

Effects of Intravascular Laser Irradiation of Blood in Mitochondria Dysfunction and Oxidative Stress in Adults with Chronic Spinal Cord Injury

Shih-Fong Huang, M.D.,¹ Yun-An Tsai, M.D.,¹ Shi-Bei Wu, Ph.D.,² Yau-Huei Wei, Ph.D.,^{2,3}
Po-Yi Tsai, M.D.,⁴ and Tien-Yow Chuang, M.D.⁴

Abstract

Objective: This study investigated the clinical effects of intravascular laser irradiation of blood (ILIB) therapy on oxidative stress and mitochondrial dysfunction in subjects with chronic spinal cord injury (SCI) resulting from trauma. **Background data:** Little is known about how ILIB may generate antioxidant defenses in humans, and there is still a lack of randomized, sham-control studies to indicate its influence on different metabolic pathways. **Methods:** Twenty-four chronic SCI subjects (assigned to a sham and a study group), and 12 normal subjects were recruited. The study group underwent 1 h daily of ILIB for 15 days over 3 weeks. The sham group underwent ILIB with no laser power. **Results:** Baseline measurements established higher oxidative stress and mitochondrial dysfunction in the SCI subjects than in the normal subjects. At day 15 of therapy, the study group revealed a significantly higher mitochondrial DNA (mtDNA) copy number, white blood cell adenosine triphosphate (WBC ATP) synthesis, and total antioxidant capacity (TAC) with significantly reduced malondialdehyde (MDA), than did the sham group. The study group intragroup comparison revealed significantly increased mtDNA copy numbers, WBC ATP synthesis, and TAC, with significantly reduced MDA, compared with its baseline measurements. The sham group intragroup comparisons demonstrated no statistical differences. Low-density lipoprotein (LDL) in the study group was significantly reduced at days 10 and 15, with significantly higher high-density lipoprotein (HDL) at day 45. **Conclusions:** Our study results contribute to the knowledge about the effectiveness of ILIB in alleviating oxidative stress and mitochondrial dysfunction in chronic SCI patients.

Introduction

INTRAVASCULAR LASER IRRADIATION OF BLOOD (ILIB) is still at the proof of concept stage; however, there are a growing number of potential clinical applications in the field of biomedicine that have been making effective use of ILIB in various pathologic processes such as pain treatment, tissue repair, and postendovascular restenosis.^{1–3} The literature to date on ILIB has documented morphologic alteration of mitochondria as a result of ILIB, as well as the activation of metabolic energy processes.^{4–8} The process of ILIB appears to modulate redox signaling in the respiratory chain through the stimulation of mitochondrial and membrane components. The production of adenosine triphosphate (ATP) and the reduction in the generation of free radicals would therefore be increased by anything that improves the flow of electrons through the respiratory chain.^{5,6}

Mitochondria are a major source of chronic reactive oxygen species (ROS), which impair lipids, protein, and mitochondria DNA, leading to the dysregulation of mitochondrial function.^{9–12} Research has indicated that mitochondrial damage may be central to the impairment of the cells of the regulatory systems – such as the nervous, endocrine and immune systems – and to the communication between them.^{13–15} Therefore, insulin-resistant type 2 diabetes mellitus, obesity, hyperglycemia, hyperlipidemia, and atherosclerosis, which are prevalent in longstanding spinal cord injury (SCI)^{16,17}, may be linked to mitochondrial dysfunction.

Although oxidative stress clearly plays an important role in the progression of SCI, most of the experiments conducted to date on oxidative stress and antioxidant defenses occurring at the injury site have been on animal models,^{18–20} and there is still a scarcity of research on mitochondrial function,

¹Department of Neuroregeneration and Neurosurgery, Taipei Veterans General Hospital, Taipei, Taiwan.

²Department of Biochemistry and Molecular Biology, National Yang-Ming University, Taipei, Taiwan.

³Department of Medicine, Mackay Medical College, New Taipei City, Taiwan.

⁴Department of Physical Medicine and Rehabilitation, Taipei Veterans General Hospital, Taipei Taiwan.

oxidative stress, and antioxidant intervention on human serum models. Moreover, although there is a considerable body of experimental data from research on humans that demonstrates the positive effects of ILIB on biostimulation, there is still a lack of randomized, sham-control studies to indicate its influence on different metabolic pathways.

Given that the emerging potential applicability of ILIB to biomedical pathologic processes has some way to go in being rigorously trialed with human subjects using appropriate methodologies, this study is a key piece of work in testing some of these in-principle clinical applications. In consideration of these, the rationale and methodology of this study were aimed at filling some of the gaps in the existing research: specifically, by characterizing temporal changes, through a single-blinded, randomized controlled methodology to investigate the effects of ILIB on mitochondrial behavior, oxidative stress biomarkers, and serum biochemical changes in chronic SCI human subjects; and, more broadly, by contributing to the body of knowledge of ILIB and its potential therapeutic effectiveness.

Methods

Participants

Thirty-six individuals were included in the study, of whom 12 were healthy controls (7 males and 5 females) and 24 (18 males and 6 females) had chronic SCI resulting from trauma, and were inpatients or outpatients of a regional SCI rehabilitation program. The SCI subjects submitted to a physical examination and routine biochemical and hematologic laboratory tests, including lipid profile, glucose, urea, creatinine, enzymes aspartate aminotransferase (AST), alanine aminotransferase, and hemograms (red blood cell [RBC], WBC, and platelet counts, and hemoglobin levels), and were receiving standard treatment, including rehabilitation therapy tailored to their specific needs.

Admission eligibility for the study included having an American Spinal Injury Association (ASIA) impairment scale classification of A, B, C, or D, a neurologic injury level between the C3 and L2 spinal cord levels, and medical stability. Subjects with communicable diseases, or active health problems (malaise, flu, fever, cough, change in urine status), or who were taking anti-inflammatory medications at the time of the study, were excluded. The subjects' clinical and demographic profiles are listed in Table 1. Prior to participation, all individuals signed an informed consent form. The investigation was in accordance with the Declaration of Helsinki Principles and conducted under the supervision of the local institutional review board protocols for human subjects and Occupational Health and Safety Administration regulations.

Allocation procedures

To achieve our primary research goal, the SCI participants were randomized into either a study or a sham group using a computer-generated random order. The randomization list was supervised by one of the investigators who had no contact with the participants until the group assignment was revealed. Further, the group assignment remained unrevealed to our laboratory staff throughout the experimental period.

TABLE 1. CLINICAL AND DEMOGRAPHIC PROFILES OF PARTICIPANTS

| Group | Gender | Age | Time post-injury(month) | Injury level | ASIA Impairment Scale |
|--------------|--------|-----|-------------------------|--------------|-----------------------|
| Study | | | | | |
| 1 | M | 57 | 20 | C4 | B |
| 2 | M | 38 | 49 | C3 | A |
| 3 | M | 29 | 6 | C6 | A |
| 4 | M | 33 | 32 | T11 | A |
| 5 | M | 37 | 136 | L2 | B |
| 6 | F | 49 | 14 | T11 | B |
| 7 | M | 48 | 74 | T3 | A |
| 8 | F | 59 | 6 | C4 | D |
| 9 | F | 53 | 28 | C4 | D |
| 10 | M | 28 | 2 | C7 | D |
| 11 | M | 63 | 7 | C4 | D |
| 12 | M | 58 | 27 | C5 | D |
| Sham | | | | | |
| 1 | M | 40 | 48 | C6 | A |
| 2 | M | 57 | 44 | C4 | D |
| 3 | M | 52 | 47 | C5 | D |
| 4 | F | 36 | 43 | T12 | B |
| 5 | F | 51 | 34 | C2 | B |
| 6 | M | 65 | 110 | L1 | D |
| 7 | M | 52 | 15 | L1 | A |
| 8 | F | 45 | 13 | C4 | A |
| 9 | M | 30 | 60 | C3 | B |
| 10 | M | 36 | 12 | T2 | A |
| 11 | M | 43 | 7 | T11 | A |
| 12 | M | 33 | 8 | C4 | A |

ASIA, American Spinal Cord Injury Association.

Intravascular low-level laser irradiation

We adopted the helium-neon (He-Ne) laser (YJ-ILIB-5, Bio-ILIB Human Energy Ltd, Taiwan) as the conditioning protocol in this study. Each participant sat on a semi-inclined sofa. A 24-gauge intravenous catheter was used for an elbow venous puncture (phlebotomy). The intravenous needle was then replaced by a fiberoptic needle inserted into the inner cannula of the intravenous catheter. The other side of the fiberoptic needle was connected to the He-Ne laser illuminator. The irradiation parameters involved in the ILIB application are shown in Table 2. For the sham control subjects, ILIB was provided without any laser power (0.0 mW). In all other respects, the program was identical for both SCI groups.

Biochemical analysis

Blood samples were taken from the SCI study group for biochemical analysis after a 12 h overnight fast at the baseline, the 10th, and the 15th days during the ILIB intervention period, and then after the end of the ILIB intervention, on the 30th day (the 45th day after the initiation of ILIB).

Isolation of WBCs from peripheral blood

Each participant provided a sample of fresh peripheral heparinized venous blood, which was centrifuged at room temperature at 3000 g for 10 min. We then transferred the buffy coat layer to 1.5 mL of Eppendorf, and washed it with RBC lysis buffer (containing 155 mM NH₄Cl, 10 mM KHCO₃, and 0.1 mM EDTA, pH 7.4) and incubated it at room

TABLE 2. PARAMETERS INVOLVED IN DETERMINING THE ILIB DOSE

| Irradiation parameter | Measurement | Irradiation time or energy delivered |
|---|---------------------------|--|
| | | Comment |
| Wavelength | 632.8 nm | Visible red low-level laser |
| Power output | 4 mW | Measured at the tip of the fiberoptic needle by optical power and energy meter |
| Irradiance (intensity or power intensity) | 2.04 W/cm ² | Irradiance = Power (W)/area (cm ²), the diameter of optical fiber aperture = 0.5 mm; area = 3.14 × (0.025 × 0.025) = 0.00196 cm ² ; irradiance = 0.004/0.00196 = 2.04 W/cm ² |
| Energy | 14.4 J | Energy(joules) = Power (W) × Time(sec) = 0.004 (W) × 3600 (sec) = 14.4 J |
| Energy density (fluence) | 7346.93 J/cm ² | Fluence = Power × time/area = 0.004(W) × 3600 (sec)/0.00196 cm ² = 7346.93 J/cm ² |
| Irradiation time | 3600 sec | One hour per each treatment |
| Treatment interval | 15 times | Fifteen times over 3 weeks |

ILIB, intravascular laser irradiation of blood.

temperature for 10 min. Each sample was centrifuged at 1500 g for 5 min at room temperature, and the cell pellets were resuspended in 1 mL of phosphate-buffered saline (PBS, pH 7.4) in preparation for the subsequent experiments. A hemocytometer was used to count the WBCs. The total number of viable WBCs was determined according to their exclusion of 0.4% Trypan Blue (Sigma-Aldrich).

Measurement of mitochondrial DNA (mtDNA) copy number from leukocytes

To measure the copy number of the mtDNA in the leukocytes, we followed a real-time polymerase chain reaction (PCR) technique according to a method described earlier by Lee et al.²¹, using a LightCycler-FastStart DNA Master SYBR Green I Kit. An aliquot of 50 ng of DNA was amplified in a 10 μ L reaction capillary with 0.25 μ L of each primer (20 μ M), 5 μ L of LightCycler SyBr Green mixed reagent and 3.5 μ L of PCR-grade H₂O. We established the real-time PCR condition as: 10 min start at 95°C, a subsequent 45 cycles of 20 sec each at 95°C; 20 sec at 62°C; and 20 sec at 72°C. At the end of every extension phase, we measured the fluorescence intensity of the PCR product at 79°C. The sequences of primers used to amplify the mitochondrial DNA (mitochondrial NADH dehydrogenase 1 gene, ND1) were: ND1R, 5'-ACCATTTGCA GACGCCATAA-3' and ND1F, 5'-TGAAATTGTTGGGCT ACGG-3'. The sequences of primers used to amplify the nuclear DNA (18S rRNA) were: 18S1546F, 5'-TAGAGGGA CAAGTGGCGTTC-3' and 18S1650R, 5'-CGCTGAGCCAGT CAGTGT-3'. We normalized the crossing points (Cp) in quantitative PCR curves between the ND1 and 18S rRNA genes using RelQuant software (Roche Applied Sciences, Mannheim, Germany) to measure the relative mtDNA copy number. To check the individual experiments for the quantification of mtDNA copy numbers performed by RT-PCR, we used DNA from CCD966SK normal skin fibroblasts as the internal standard for calibrating the relative mtDNA copy numbers in each sample from each individual RT-PCR run. The Ct (ND1 to 18S) for CCD966SK was therefore defined as 1.0, and we were therefore able to calculate the relative mtDNA copy number for each sample in each individual experiment set. In this way, we precluded any errors from arising from an individual quantitative PCR run. The PCR product for ND1 is 108 base pairs and for 18S, 105 base pairs. The amplification of the PCR product was followed by a melting curve analysis to monitor the homogeneity of the

amplicons. At the final RT-PCR step, melting curve analysis was performed, consisting of a denaturation at 95°C followed by a renaturation step by a slow melt approaching 68°C with a ramping rate of 0.2°C/sec with continuous data acquisition.

Measurement of total cellular ATP in WBCs

We measured the total cellular ATP content of the WBCs using the Bioluminescent Somatic Cell Assay Kit (Sigma-Aldrich, St. Louis, MO) in accordance with the manufacturer's instructions and our experience.²² Once the WBCs were separated from the whole blood samples, we released the total cellular ATP contents by mixing an aliquot of 50:1 WBC suspension with 150:1 Somatic Cell Releasing Reagent. One half (100:1) of the mixture was then transferred into a black OptiPlateTM-96F 96-well plate (Packard Biosciences, Groningen, The Netherlands) containing 100:1 ATP Assay Mix. We then measured the luminescence intensity using the Victor²_{TM} 1420 multilabel counter machine (PerkinElmer Life Sciences, Inc., Boston, MA). The ATP sample standards in the range of 0–300 pmoles were used, and each ATP value was normalized with WBC numbers.

Analysis of malondialdehyde (MDA) in blood serum

MDA in the participants' blood serum was measured using an MDA assay kit (MDA-586, OxisResearch, Inc. Portland, OR) with spectrophotometric analysis. Measurements were made according to the manufacturer's instruction and our own experience.²³ The assay was involved in the reaction with a chromogenic reagent N-methyl-2-phenylindole (NMPI) to form an intensely colored carbocyanine dye with a maximum absorption of 586 nm. The methodology we used here is specific to MDA, rather than to other lipid peroxidation byproducts such as 4-hydroxy alkenal, because under the same experimental conditions they are unable to produce significant absorbance at 586 nm. Using this assay, an MDA standard curve was established at the concentration range of 0–50 μ M.

Determination of total antioxidant capacity (TAC) of blood serum

The TAC of the participants' serum was measured using a PAO kit (Japan Institute for the Control of Aging, Nikken SEIL Co., Fukuroi, Japan) in accordance with the manufacturer's instruction and our own experience.²² An aliquot of

100 μ L of diluted serum from each participant was mixed with 50 μ L of Cu^{2+} solution, which can be reduced to form Cu^+ in correspondence with the antioxidant status of serum after 5 min of incubation at room temperature. We then added an aliquot of 50 μ L of a chromogenic solution to react with the Cu^+ ion. The absorbance at 490 nm was measured immediately. We determined the TAC levels of the serum by reducing the Cu^{2+} solution with corresponding concentrations of uric acid. To gauge the antioxidant power of the samples, we multiplied the corresponding uric acid concentration by 2189 (1 mM uric acid = 2189 μ mole/L copper reducing power).

Statistical analysis

Statistical analysis of the processed data was performed with commercial software (SPSS 17.0 for Windows; SPSS Inc.). Descriptive statistics were used to elucidate a set of known data in a clear and concise manner, as in term of its mean and variance. Herein we used descriptive statistics to summarize a data set of subject characteristics and display major findings. The Wilcoxon signed-rank test was used to compare the biochemistry changes across time versus the baseline values of the study group. One-way analysis of variance (ANOVA) was used to analyze the baseline data of MDA, TAC, mtDNA and total cellular ATP in WBC among the normal, sham, and study groups. A two-way repeated measures ANOVA was used to analyze antioxidant activity of the body and mitochondrial activity with time (baseline, day 1, and day 15 post-intervention) as the within- and between-group factor. When an intervention between group and time was found, *post-hoc* tests were performed to determine the real within- and between- group differences. The level of significance was set at 0.05.

Results

Across all demographic indicators, there were no significant differences between the sham and study groups (Table 1).

The complete blood cell counts, liver transaminases, blood urea nitrogen, creatinine, creatinine phosphokinase, and urine pH were unchanged before and after the intervention. However, low-density lipoprotein (LDL) in the study group showed a significant decrease at both days 10 and 15, with significantly higher high-density lipoprotein (HDL) found at day 45 as compared with the baseline measurement (Table 3).

The baseline measurements revealed that there were significant differences among groups. The normal, aged-matched subjects has a significantly higher mtDNA copy number, WBC ATP synthesis, and TAC, and less MDA than did the study and sham subjects of the SCI group (Fig. 1). At day 15 of the ILIB intervention, the study group was found to have a significantly higher mtDNA copy number, WBC ATP synthesis, and TAC with significantly reduced MDA than did the sham group (Fig. 2). The intragroup comparisons for the study group revealed that the mtDNA copy number, WBC ATP synthesis, and TAC at day 15 were significantly increased, with significantly reduced MDA (Fig. 2). The intragroup comparisons for the sham group, by contrast, demonstrated no statistical differences for any biomarkers.

Discussion

Use of He-Ne laser (632.8 nm) to promote functional regeneration of mitochondria and pro-oxidant/antioxidant equilibrium in living bodies is common; moreover ILIB with He-Ne laser has been applied in aid of treatment of acute cerebral infarction, rheumatoid arthritis, arteriosclerosis of the leg, and redifferentiation therapy of malignant tumors.²⁴⁻²⁸ However, over the past 30 years, there has been a small, but growing, body of clinical studies that has been challenging the status-quo belief with regard to the therapeutic effectiveness of ILIB, and at present there are no generally accepted practical ILIB strategies. Expectations that ILIB can reduce oxidative stress in chronic SCI patients have to date

TABLE 3. THE SERUM BIOMARKERS AND PHYSICAL EXAMINATION CHANGES IN STUDY GROUP BEFORE AND AFTER ILIB INTERVENTION

| | Baseline | Day 10 | Day 15 | Day 45 |
|------------------------------|------------------|------------------------------|------------------------------|----------------------------|
| | Mean (SD) | Mean (SD) | Mean (SD) | Mean (SD) |
| Uric acid | 6.88 (1.904) | 6.34 (1.896) | 6.42 (1.733) | 7.21 (2.549) |
| Cholesterol | 195.42 (31.070) | 189.75 (36.834) | 182.00 (36.287) | 200.00 (30.887) |
| TG | 221.58 (158.882) | 246.08 (171.684) | 235.08 (187.117) | 216.92 (140.864) |
| HDL | 22.33 (7.981) | 23.92 (6.459) | 22.75 (6.811) | 30.00 (6.523) ^a |
| LDL | 132.25 (28.566) | 117.58 (34.140) ^a | 114.17 (31.287) ^a | 128.58 (38.451) |
| Glucose AC | 114.08 (42.361) | 107.17 (31.963) | 110.58 (32.779) | 111.17 (40.994) |
| Insulin AC | 10.48 (8.861) | 11.86 (8.466) | 10.80 (11.116) | 10.50 (11.142) |
| IL-6 | 8.08 (4.544) | 7.34 (1.897) | 9.75 (4.078) | 9.23 (2.931) |
| TNF- α | 13.08 (15.226) | 12.43 (13.145) | 11.26 (12.298) | 12.64 (14.695) |
| Adiponectin | 6.84 (5.675) | 7.26 (7.077) | 6.92 (6.618) | 6.97 (7.907) |
| Abdominal circumference (cm) | 93.6 (9.73) | 93.3 (9.99) | 93.1 (9.79) | 93.2 (11.66) |
| Systolic pressure (mm Hg) | 120 (18.0) | 114 (17.2) | 115 (14.2) | 110 (18.9) |
| Diastolic pressure (mm Hg) | 83 (13.5) | 76 (8.8) | 77 (11.2) | 75 (9.6) |
| Body weight (kg) | 75.7 (14.9) | 75 (14.3) | 75 (15.2) | 75 (15.3) |

Values were Mean \pm SD.

Wilcoxon signed-rank test was used to compare the data changes across time versus baseline in study group.

^a*p* < 0.05.

ILIB, intravascular laser irradiation of blood; TG, triglycerides; HDL, high-density lipoprotein; LDL, low-density lipoprotein; IL, interleukin; TNF, tumor necrosis factor.




FIG. 1. (A) The baseline comparisons of mitochondrial DNA copy number among normal, study, and sham groups. The normal group had significant higher mitochondrial copy number than did the study and sham groups ($*p < 0.05$). (B) The baseline comparisons of white blood cell (WBC) intracellular adenosine triphosphate (ATP) among normal, study, and sham groups. The normal group had significant higher WBC intracellular ATP than did the study and sham groups ($*p < 0.05$). (C) The baseline comparisons of serum malondialdehyde (MDA) among normal, study, and sham control groups. *The normal group had significant lower serum MDA than did the study and sham groups ($*p < 0.05$). (D) The baseline comparisons of serum total antioxidative capacity (TAC) among normal, study, and sham groups. The normal group had significant higher serum TAC than did the study and sham groups ($*p < 0.05$).

been based on findings that mitochondrial photostimulation enhances ATP synthesis and contributes to redox signaling in the clearing of free radicals;²⁹ and on the assumption that mitochondria are the primary targets of red light radiation with 632.8nm wavelength of whole cells.^{5-7,29} This study is the first to rigorously characterize temporal changes in mitochondrial function and antioxidant activity following ILIB therapy in chronic SCI subjects. Although the overall procedure required 15 days of 1h 4mW He-Ne laser ILIB intervention sessions, all of the subjects were able to tolerate the whole procedure with no adverse effects, which is consistent with and reinforces previous findings.⁴

The main findings of this study were: (1) that chronic SCI patients were indisputably in states of mitochondrial damage and oxidative stress in comparison with normal subjects;

(2) there was a significant increase in the mtDNA copy number, WBC ATP synthesis, and TAC, with significantly reduced MDA production in the study group as compared with the sham group at day 15 of ILIB intervention; 3) there was a significant increase in the mtDNA copy number, WBC ATP synthesis, and TAC, with significantly reduced MDA production in the study group participants at day 15 of ILIB intervention as compared with their baseline values; and 4) there was significant decrease in LDL of the study group participants at day 10 and day 15 of ILIB intervention, with significant increase in their HDL as compared with their baseline measurements.

In vivo data are still limited concerning the mechanism by which secondary SCI deterioration occurs following initial primary injury. However, several lines of evidence have




FIG. 2. (A) The inter- and intragroup comparisons of mitochondrial DNA copy numbers in study and sham groups. Significantly higher mitochondrial DNA copy numbers were noted at day 15 in the study group compared with baseline (intragroup) or sham group (intergroup). 1, baseline; 2, day 1 post-intervention; 3, day 15 post-intervention. (B) The inter- and intragroup comparisons of white blood cell adenosine triphosphate (WBC ATP) in study and sham control groups. Significantly higher WBC ATPs were noted at day 15 in study group compared with baseline (intragroup) or sham group (intergroup). 1, baseline; 2, day 1 post-intervention; 3, day 15 post-intervention. (C) The inter- and intragroup comparisons of serum malondialdehyde (MDA) in study and sham control groups. Significantly lower serum was noted at day 15 in the study group compared with baseline (intragroup) or sham group (intergroup). 1, baseline; 2, day 1 post-intervention; 3, day 15 post-intervention. (D) The inter- and intragroup comparisons of serum TAC in study and sham control groups. Significantly higher serum total antioxidative capacity (TAC) was noted at day 15 in the study group compared with baseline (intragroup) or sham group (intergroup). 1, baseline; 2, day 1 post-intervention; 3, day 15 post-intervention.

suggested that ROS formation and the consequent development of oxidative stress are important events associated with SCI.²⁰ ROS are difficult to measure directly as they are highly reactive and short lived. Indirect measures such as MDA and other biomarkers are therefore generally used to predict the amount of ROS or the extent of oxidative stress occurring in the SCI subjects.³⁰ The baseline findings of this study, which established that chronic SCI subjects had significantly higher baseline MDA content, lower TAC, and impaired mitochondrial metabolic activities than did normal subjects, support the abovementioned evidence of the development of ROS formation and oxidative stress in secondary SCI deterioration. The data recorded for the chronic SCI study and sham groups of this research also confirm previous findings that the application of a He-Ne laser could improve mito-

chondrial activities (DNA and ATP synthesis) in WBC and reinforce antioxidant defenses in the blood serum of chronic SCI patients. Although the underlying mechanisms have yet to be clearly elucidated, our findings do point to a potential link between photostimulation and mitochondrial proliferation and/or cellular homeostasis. As ILIB is known to generate low levels of ROS that can upregulate cellular antioxidant defenses and improve cell signaling,³¹ it is feasible that TAC increases as a result of ILIB intervention in compensation for a beneficial, low-level increase in cellular ROS, and that serum MDA decreases because the proximate membrane protein-glutathione peroxidase 4, which is responsible for lipid peroxidation, is downregulated.³² Our results here support the concept that ROS act as key secondary messengers contributing to ILIB-induced biologic

outcomes. As the prevalence of adiposity soars in chronic SCI patients, appreciation of the close links between obesity and disease increases. The strong relationships between lipid metabolism and poor health outcomes, including cardiovascular disease, diabetes, and metabolic syndrome, mandate clarification of cholesterol constituent changes in chronic SCI patients. It is of interest to note that the LDL of the study group decreased during the application period of ILIB and that its HDL was subsequently raised. The redistribution of cholesterol was strongly evidenced in concordance with markedly reducing serum MDA and increasing serum TAC. We consequently propose that ILIB plays a central role in the regulation of lipid metabolism, which improves LDL catabolism, reduces the oxidation of LDL, and increases HDL levels. This beneficial effect of ILIB on LDL catabolism could be partially related to the upregulation of LDL receptor expression.^{33–35} The mechanism by which ILIB increases HDL is not fully understood, but is likely related to increased expression of lipoprotein lipase or decreases in apolipoprotein (Apo) A-I clearance,^{34,36} which may augment the biosynthesis of HDL.³⁴ It is clear that *in vivo*, HDL can inhibit the oxidative modification of LDL. In addition to the antioxidative Apo A-I, glutathione and paraoxonase 1 are all HDL-associated enzymes, which inactivate lipid peroxides and other oxidized LDL lipid components.^{36,37}

Limitations

There were several limitations to the methodology of this study. The lack of continuous monitoring of mitochondrial function and of antioxidant effects before and after each ILIB session made it impossible to examine the precise moment at which mitochondrial proliferation and cellular homeostasis were reached and their duration. However, our decision to administer ILIB 15 times in this protocol was in accordance with previous studies and suggestions by Weber et al.⁴ Further research on different laser energies and application settings would advance these findings and identify optimal treatments for adults with chronic SCI. This study focused on the proof that oxidative stress in chronic SCI subjects is alleviated after exposure to He-Ne laser. The relationship between the generation of antioxidant defenses and clinical motor/sensory improvement was beyond the scope of this study comprising a small number of participants undergoing a short-term period of treatment. The achievement of significance for every study group participant of this study firmly indicates the value of further exploring and testing its findings in a large-scale trial of ILIB intervention as part of longer-term treatment of chronic SCI patients. We expect that more alleviation of oxidative stress in longer-term trials may have better results in neurologic and metabolic recovery of SCI, as oxidative stress, in addition to ROS, is a key element in the pathophysiology of SCI.^{18–20}

Conclusions

The baseline findings of this study across normal and chronic SCI patients established that our chronic SCI patients were in states of mitochondrial damage and oxidative stress. ILIB intervention was found to significantly increase subjects' mtDNA copy number, WBC ATP synthesis, and TAC; significantly reduce MDA production; significantly decrease LDL; and significantly increase HDL. Our findings on

human subjects have therefore reinforced the body of research that contends that ILIB therapy promotes mitochondrial DNA or total cellular ATP synthesis, antioxidant activity, and the reduction of lipid peroxidation and dyslipidemia generation, contributing to the alleviation of chronic conditions, such as chronic SCI. Given this precedent and the alleviation of oxidative stress and ROS-scavenging capacity, we believe that this clearly signals ILIB as a potential addition to the current armament of treatment for those with SCI.

Author Disclosure Statement

No competing financial interests exist.

References

1. Timofeyev, V.T., Poryadin, G.V., and Goloviznin, M.V. (2001). Laser irradiation as a potential pathogenetic method for immunocorrection for rheumatoid arthritis. *Pathophysiology* 8, 35–40.
2. Arora, H., Pai, K.M., Maiya, A., Vidyasagar, M.S., and Rajeev, A. (2008). Efficacy of He-Ne laser in the prevention and treatment of radiotherapy-induced oral mucositis in oral cancer patients. *Oral Surg. Oral Med. Pathol. Oral Radiol. Endod.* 105,180–186.
3. DerKacz, A., Protasiewicz, M., Poreba, R., Szuba, A., and Andrzejak, R. (2010). Usefulness of intravascular low power laser illumination in preventing restenosis after percutaneous coronary intervention. *Am. J. Cardiol.* 106(8), 1113–1117.
4. Weber, M.H., Fußgänger-May, T., and Wolf, T. (2007). The intravenous laser blood irradiation – introduction of a new therapy. *Dt. Ztschr. F. Akup.* 50,12–23.
5. Karu, T.I., Pyatibrat, L.V., Kolyakov, S.F., and Afanasyeva, N.I. (2008). Absorption measurements of cell monolayers relevant to mechanisms of laser phototherapy: reduction or oxidation of cytochrome c oxidase under laser radiation at 632.8 nm. *Photomed. Laser Surg.* 26, 593–599.
6. Karu T.I. (2008). Mitochondrial signaling in mammalian cells activated by red and near-IR radiation. *Photochem. Photobiol.* 84, 1091–1099.
7. Hu, W.P., Wang, J.J., Yu, C.L., Lan, C.C., Chen, G.S., and Yu, H.S. (2007). Helium-neon laser irradiation stimulates cell proliferation through photostimulatory effects in mitochondria. *J. Invest. Dermatol.* 127, 2048–2057.
8. Sroka, R., Schaffer, M., Fuchs, C., et al. (1999). Effects on the mitosis of normal and tumor cells induced by light treatment of different wavelengths. *Lasers Surg. Med.* 25, 263–271.
9. Fernández-Checa, J.C., Fernández, A., Morales, A., Mari, M., García-Ruiz, C., and Colell, A. (2010). Oxidative stress and altered mitochondrial function in neurodegenerative diseases: lessons from mouse models. *CNS Neurol. Disord. Drug Targets* 9,439–454.
10. Hipkiss, A.R. (2010). Mitochondrial dysfunction, proteotoxicity, and aging: causes or effects, and the possible impact of NAD⁺-controlled protein glycation. *Adv. Clin. Chem.* 50,123–150.
11. Leduc, L., Levy, E., Bouity-Voubou, M., and Delvin, E. (2010). Fetal programming of atherosclerosis: possible role of the mitochondria. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 149, 127–130.
12. Vaishnav, R.A., Singh, I.N., Miller, D.M., and Hall, E.D. (2010). Lipid Peroxidation-Derived Reactive Aldehydes Directly and Differentially Impair Spinal Cord and Brain Mitochondrial Function. *J. Neurotrauma.* 27, 1311–1320.

13. Frost, F., Roach, M.J., Kushner, I., and Schreiber, P. (2005). Inflammatory C-reactive protein and cytokine levels in asymptomatic people with chronic spinal cord injury. *Arch. Phys. Med. Rehabil.* 86, 312–317.
14. Rajan, S., McNeely, M.J., Hammond, M., Goldstein, B., and Weaver, F. (2010). Association between obesity and diabetes mellitus in veterans with spinal cord injuries and disorders. *Am. J. Phys. Med. Rehabil.* 89, 353–361.
15. Inskip, J., Plunet, W., Ramer, L., Ramsey, J.B., Yung, A., Kozlowski, P., Ramer, M., and Krassioukov, A. (2010). Cardiometabolic risk factors in experimental spinal cord injury. *J. Neurotrauma.* 27, 275–285.
16. Mokini, Z., Marcovecchio, M.L., and Chiarelli, F. (2010). Molecular pathology of oxidative stress in diabetic angiopathy: role of mitochondrial and cellular pathways. *Diabetes Res. Clin. Pract.* 87, 313–321.
17. Liang, H., Mojtahedi, M.C., Chen, D., and Braunschweig, C.L. (2008). Elevated C-reactive protein associated with decreased high-density lipoprotein cholesterol in men with spinal cord injury. *Arch. Phys. Med. Rehabil.* 89, 36–41.
18. Vaziri, N.D., Lee, Y.S., Lin, C.Y., Lin, V.W., and Sindu, R.K. (2004). NAD(P)H oxidase, superoxide dismutase, catalase, glutathione peroxidase and nitric oxide synthase expression in subacute spinal cord injury. *Brain Res.* 995, 76–83.
19. Azbill, R.D., Mu, X.J., Bruce-Keller, A.J., Mattson, M.P., and Springer, J.E. (1997). Impaired mitochondrial function, oxidative stress and altered antioxidant enzyme activities following traumatic spinal cord injury. *Brain Res.* 765, 283–290.
20. Jia, Z., Zhu, H., Li, J., Wang, X., Misra, H., and Li, Y. (2012). Oxidative stress in spinal cord injury and antioxidant-based intervention. *Spinal Cord* 50, 264–274.
21. Lee, H.C., Li, S.H., Lin, J.C., Wu, C.C., Yeh, D.C., and Wei, Y.H. (2004). Somatic mutations in the D-loop and decrease in the copy number of mitochondrial DNA in human hepatocellular carcinoma. *Mutat. Res.* 547, 71–78.
22. Chen, C.T., Shih, Y.R., Kuo, T.K., Lee, O.K., and Wei, Y.H. (2008). Coordinated changes of mitochondrial biogenesis and antioxidant enzymes during osteogenic differentiation of human mesenchymal stem cells. *Stem Cells* 26, 960–968.
23. Chen, S.S., Wu, S.B., and Wei, Y.H. (2010). The difference in oxidative stress of the blood between using 5% glucose water and distilled water as the irrigant for BPH patients undergoing transurethral resection of the prostate. *World J. Urol.* 28, 33–37.
24. Li, X.Z., Li, M.X., and Wang, Y.F. (1998). He-Ne laser intravascular irradiation therapy in the treatment of 55 cases with acute cerebral infarction. *Chin. J. Phys. Ther.* 21, 17–19.
25. Huo, X.L., Zhang, Y.D., Di, Q., Wu, X.F., Fang, Q., and Hu, Z.R. (2000). Effect of the intravascular low level laser irradiation with different doses of different lasers on the blood biomarkers in patients with cerebral infarction. *Chin. J. Laser Med. Surg.* 9, 80–84.
26. Timofeyev, V.T., Poryadin, G.V., and Goloviznin M.V. (2001). Laser irradiation as a potential pathogenetic method for immunocorrection in rheumatoid arthritis. *Pathophysiology* 8, 35–40.
27. Czernicki, J., Radziszewski, K., and Talar, J. (1994). Effect of laser biostimulation on leg blood flow in the course of arteriosclerosis. *Pol. Tyg. Lek.* 25, 363–365.
28. Andra, F. (2007). The effects of intravascular low level laser therapy in the scope of a redifferentiation therapy of malignant tumors. *German Med. J. Oncol.* 6,1–5.
29. Tafur, J., and Mills, P.J. (2008). Low-intensity light therapy: exploring the role of redox mechanisms. *Photomed. Laser Surg.* 26, 323–328.
30. Arcari, D.P., Porto, V.B., Rodrigues, E.R.V., et al. (2011). Effect of mate tea (*Ilex paraguariensis*) supplementation on oxidative stress biomarkers and LDL oxidisability in normo- and hyperlipidaemic humans. *J. Funct. Foods* 3, 190–197.
31. Zhang, J., Xing, D., and Gao, X. (2008). Low-power laser irradiation activates Src tyrosine kinase through reactive oxygen species-mediated signaling pathway. *J. Cell Physiol.* 217, 518–528.
32. Van Raam, B.J., Sluiter, W., Wit, E., Roos, D., Verhoeven, A.J., and Kuijpers, T.W. (2008). Mitochondrial membrane potential in human neutrophils is maintained by complex III activity in the absence of supercomplex organization. *PLoS One* 3, 1–12.
33. Galland, F., Duvillard, L., Petit, J.M., et al. (2006). Effect of insulin treatment on plasma oxidized LDL/LDL-cholesterol ratio in type 2 diabetic patients. *Diabetes Metab.* 32,625–631.
34. Rezen, T., Roman, D., Pascussi, J.M., and Monostory, K. (2011). Interplay between cholesterol and drug metabolism. *Biochim Biophys Acta* 1814,146–160.
35. Alving, C.R., and Wassef, N.M. (1999). Naturally occurring antibodies to cholesterol: a new theory of LDL cholesterol metabolism. *Immunol. Today.* 20(8),362–366.
36. Link, J.J., Rohatgi, A., and De Lemos, J.A. (2007). HDL cholesterol:physiology, pathophysiology and management. *Curr. Probl. Cardiol.*32, 268–314.
37. Raveh, O., Pinchuk, I., Fainaru, M., and Lichtenberg, D. (2001). Kinetics of lipid peroxidation in mixtures of HDL and LDL mutual effects. *Free Radic. Biol. Med.* 31(11), 1486–1497.

Address correspondence to:

Tien-Yow Chuang
 Department of Physical Medicine and Rehabilitation
 National Yang-Ming University
 and Taipei Veterans General Hospital
 No. 201, Sec. 2, Shih-Pai Road
 Taipei
 Taiwan 11217

E-mail: tychuang@vghtpe.gov.tw

This article has been cited by:

1. Calheiros Andrea Paiva Corsetti, Moreira Maria Stella, Gonçalves Flávia, Aranha Ana Cecília Correa, Cunha Sandra Ribeiro, Steiner-Oliveira Carolina, Eduardo Carlos de Paula, Ramalho Karen Müller. 2017. Photobiomodulation in the Prevention of Tooth Sensitivity Caused by In-Office Dental Bleaching. A Randomized Placebo Preliminary Study. *Photomedicine and Laser Surgery* 35:8, 415-420. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
2. Jenna L. Gollihue, Alexander G. Rabchevsky. 2017. Prospects for therapeutic mitochondrial transplantation. *Mitochondrion* 35, 70-79. [[Crossref](#)]
3. Fernanda Rossi Paolillo, Audrey Borghi-Silva, Ross Arena, Nivaldo Antonio Parizotto, Cristina Kurachi, Vanderlei Salvador Bagnato. 2017. Effects of phototherapy plus physical training on metabolic profile and quality of life in postmenopausal women. *Journal of Cosmetic and Laser Therapy* 28, 1-9. [[Crossref](#)]
4. Wan-Hua Yang, Shiou-Ping Lin, Shin-Tsu Chang. 2017. Case report. *Medicine* 96:2, e5646. [[Crossref](#)]
5. Cíntia Yuki Fukuoka, Gabriella Torres Schröter, José Nicolau, Alyne Simões. 2016. Low-power laser irradiation in salivary glands reduces glycemia in streptozotocin-induced diabetic female rats. *Journal of Biophotonics* 9:11-12, 1246-1254. [[Crossref](#)]
6. Takahiro Ando, Michael R. Hamblin. Chapter 23 Low-Level Laser Therapy for Spinal Cord Repair 415-434. [[Crossref](#)]
7. Daiane Thais Meneguzzo, Leila Soares Ferreira, Eduardo Machado de Carvalho, Cássia Fukuda Nakashima. Chapter 46 Intravascular Laser Irradiation of Blood 933-952. [[Crossref](#)]
8. Sungkyoo Lim, Jun Ho Lee, Ji Geun Jang. 2015. Red/Infrared LED Unit with Moderate Light Intensity and Low Heat Generation for Personal Phototherapy Applications. *Molecular Crystals and Liquid Crystals* 618:1, 1-9. [[Crossref](#)]
9. Efremova Yulia, Sinkorova Zuzana, Navratil Leos. 2015. Protective Effect of 940nm Laser on Gamma-Irradiated Mice. *Photomedicine and Laser Surgery* 33:2, 82-91. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
10. Bridgette D. Semple. 2014. Early preservation of mitochondrial bioenergetics supports both structural and functional recovery after neurotrauma. *Experimental Neurology* 261, 291-297. [[Crossref](#)]
11. Samir P. Patel, Patrick G. Sullivan, Jignesh D. Pandya, Glenn A. Goldstein, Jenna L. VanRooyen, Heather M. Yonutas, Khalid C. Eldahan, Johnny Morehouse, David S.K. Magnuson, Alexander G. Rabchevsky. 2014. N-acetylcysteine amide preserves mitochondrial bioenergetics and improves functional recovery following spinal trauma. *Experimental Neurology* 257, 95-105. [[Crossref](#)]
12. Yong Ye, Yue Li, Fei Fang. 2014. Opening of brain blood barrier induced by red light and central analgesic improvement of cobra neurotoxin. *Journal of Photochemistry and Photobiology B: Biology* 134, 16-22. [[Crossref](#)]
13. Francisco J. Ordonez, Miguel A. Rosety, Alejandra Camacho, Ignacio Rosety, Antonio J. Diaz, Gabriel Fornieles, Marco Bernardi, Manuel Rosety-Rodriguez. 2013. Arm-Cranking Exercise Reduced Oxidative Damage in Adults With Chronic Spinal Cord Injury. *Archives of Physical Medicine and Rehabilitation* 94:12, 2336-2341. [[Crossref](#)]
14. Dragos Andrei Chiran, Gerhard Litscher, Michael Weber, Laura Marinela Ailioaie, Constantin Ailioaie, Daniela Litscher. 2013. Intravenous Laser Blood Irradiation Increases Efficacy of Etanercept in Selected Subtypes of Juvenile Idiopathic Arthritis: An Innovative Clinical Research Approach. *Evidence-Based Complementary and Alternative Medicine* 2013, 1-9. [[Crossref](#)]