# **I-131 Ablation Therapy for Cancer**

(4<sup>th</sup> version Aug.1, 2020)

## **Patient Preparation:**

- 1. Injection of contrast media in radiological examination such as CT scan, IVP test and Bronchography should be avoided in at least one months before the test.
- 2. If you have any possibility for pregnancy, please tell us in advance.
- **3.** Ten days before the test, please start low-iodine diet including low-iodine salt. Don't take sea food, kelp, eggs, chocolate, and any drug containing iodine such as multivitamin, coughing drug, etc.
- **4.** Discontinue thyroid hormone for at least four weeks before the treatment. Other drugs depend on individual circumstance. Please consult doctors.
- 5. If you are prescribed to take doses more than 30 mCi, you will be admitted to the isolation ward designed for radiation protection.

### **Before Treatment:**

- 1. Check serum TSH and thyroglobulin before scanning if necessary
- 2. Follow the instruction of pharmacists. Take a capsule or solution containing I-131.

#### **After Treatment:**

- **1.** Fasting for 4 hours after intake of I-131.
- 2. In order to reduce radiation burden to non-target organs, suggest you drink more water and empty bladder frequently. If constipation, laxatives is suggested.
- **3.** In the first day post I-131 intake, liquid or semi-liquid meal is suggested to reduce the probability of nausea and vomiting.
- **4.** Two days after intake of the I-131, you can take drugs (including thyroid hormone) or food as you need.
- 5. Within 7 days post I-131 intake, please frequently consume a hard & sour candy to stimulate the secretion of saliva, which can reduce I-131 accumulation in salivary glands.
- 6. About one week after I-131 intake, a whole body scan will be performed. Some patients may need to take the I-131 more than one time.
- 7. Within seven days post I-131 intake, please try not to hold children or stay with people for a long time (over half hour) and please sleep alone.
- **8.** I-131 can excrete via lactation. Please stop breast feeding for at least one week after I-131 intake.

#### Follow-Up:

After iodine-131 cancer eradication treatment, our medical staff will assign a whole body scan for you. Remember to return to the hospital as scheduled.

#### Please feel free to contact us if you have any question.

TEL: 04-23592525 extend 4801 Email: <u>m@vghtc.gov.tw</u>

Nuclear Medicine Division Taichung Veterans General Hospital