

臺中榮總 兒科適用水劑藥品清單

藥品成份/商品名	成份含量	一般劑量
<u>Analgesics/Antipyretics (止痛退燒藥)</u>		
* Acetaminophen Anti-phen®syrup 安佳熱糖漿	24 mg/ml, 60ml/bot	Fever: 10-15 mg/kg/dose PO Q4H-Q6H, maximum 5 doses/day
Ibuprofen Ibuprofen oral® susp 依普芬口服懸液劑	20 mg/ml, 60 ml/bot	6 m/o-12 yrs: 0.25-0.5 ml/kg
<u>Epectorants (祛痰劑)</u>		
Glyceryl guaiacolate Guaphen® Syrup 120mL 咳酚糖漿	20 mg/ml, 120 ml/bot	一天4~6次，每次間隔至少4小時。 成人及12歲以上：10毫升 6歲以上未滿12歲：5毫升 3歲以上未滿6歲：2.5毫升 3歲以下：請洽醫師診治
<u>Mucolytic Drugs (黏液溶解劑)</u>		
* Bromhexine Bisolvon® solution 氣舒痰液	2 mg/ml, 50 ml/bot	Infant : 0.5ml TID 5 yrs : 1ml TID 5-10 yrs: 2 ml TID
<u>Antitussives (鎮咳劑)</u>		
Butamirate citrate Sutussi®liquid 停咳喜液	0.8 mg/mL, 120 mL/bot	Adults : 20-25 mg QID ≥ 12 years : 15-22 mg QID 6-12 years : 6-15 mg (7.5-15 ml) QID 3-6 years : 5-6 mg (7.5 ml) QID 1-3 years : 3 mg (3.75 mL) QID 2 month-1 years : 2 mg (2.5 ml) QID
<u>Antihistamines (抗組織胺藥)</u>		
* Cetirizine HCl Cetirizine®Sol'n 勝克敏液	1 mg/ml, 60 ml/bot	3-6 y/o: 5 ml/day QD~BID >6 y/o: 10 ml QD
* Cyproheptadine HCl Cyproh®syrup 喜普液	0.4 mg /ml, 60 ml/bot	2-6 yrs: 5 ml TID 6-14yrs : 10 ml TID
Ketotifen Athmin®syrup 喘敏糖漿	0.2 mg/ml, 60ml/bot	0.5-2 mg BID
<u>Autonomic-β-Adrenergic Agonists</u>		
* Procaterol Meptin®Liquid 滅喘淨液	5 mcg/ml, 60 ml/bot	< 6 y/o: 10-15 mcg QD > 6 y/o: 25 mcg QD-BID

臺中榮總 兒科適用水劑藥品清單

藥品成份/商品名	成份含量	一般劑量
Respir. Smooth Muscle Relax. (呼吸系統平滑肌鬆弛劑)		
* Theophylline Centertheo®liquid 適優喘液	5.34 mg/ml, 60 ml/bot	6-12 yrs: 125-250 mg Q6H (3 ml/kg/day Q8-6H or 1 ml/kg/dose)
Misc. Respiratory Tract Agents (其他呼吸道治療劑)		
Chlorpheniramine + Methylephedrine + guaiaconoL Secorine®syrup 息咳寧糖漿	Each ml: Chlorpheniramine 0.1mg + Methylephedrine 1mg + guaiaconoL glycolate 5 mg. 120mL/bot	Adult : 20 mL 3~4 times a day. 9-12 yrs: 10 mL 3~4 times a day. 6-9 yrs: 6.7 mL 3~4 times a day.
Antibacterials (抗生素)		
* Amoxicillin+Potassium clavulanate Curam®oral syrup 諾快寧口服懸液用粉	Each 5ml: Amoxicillin 250 mg + Potassium clavulanate 62.5mg 60 ml/bot	● 2-12 yrs: 30~60mg/kg/day Q8H < 2 yrs: 30~40mg/kg/day Q8H (Amoxicillin dose) ● 須先加水至瓶身凹陷標記處，並充分振搖溶解，每次使用前也需搖勻。 ● 稀釋後需冷藏，安定性為7天。
* Azithromycin Zithromax®susp 日舒懸液用粉	200 mg/5 ml, 15 ml/bot	● <15 kg: 10 mg/kg for 3 days 15-25 kg: 200mg (5ml)/QD for 3day ● 須先加9ml水，稀釋後為15ml，並充分振搖溶解，每次使用前也需搖勻。 ● 稀釋後室溫安定性為7天。
Cephalexin Ulexin®susp 60mL 優力黴素口服懸液	25mg/mL, 1.5gm/60mL/bot	● Mild to moderate infection: 25-50 mg/kg/day Q6-12H; MAX daily dose: 2000 mg/day. ● Severe infection (eg, bone and joint infections): 75-100 mg/kg/day Q6-8H; MAX daily dose: 4000 mg/day.
Antifungals(抗黴菌)		
* Nystatin Mycostatin®susp 滅菌靈懸液用粉劑	100,000 U/ml, 24 ml/bot	● 1 ml QID (需要時可增加劑量) ● 須先加23ml水稀釋，並充分振搖溶解，每次使用前也需搖勻。 ● 稀釋後室溫安定性為7天。
Anticonvulsants (抗痙攣劑)		
Cannabidiol Epidyolex® soln 100mg/mL	100 mg/mL, 100mL/bot	● Dravet syndrome (DS) or Lennox-Gastaut syndrome (LGS) : Initial: 2.5 mg/kg BID, maintenance: 5 mg/kg BID, MAX: 20 mg/kg/day. ● Tuberous sclerosis complex (TSC): Initial: 2.5 mg/kg BID, maintenance: 12.5 mg/kg BID, MAX: 25 mg/kg/day.

臺中榮總 兒科適用水劑藥品清單

藥品成份/商品名	成份含量	一般劑量
* Levetiracetam Keppra® oral solution 優閒內服液劑	100 mg/ml, 300 ml/bottle	4-15 years: 10 mg/kg/dose BID, may increase every 2 weeks by 10 mg/kg/dose to a maximum of 30 mg/kg/dose BID.
* Oxcarbazepine Trileptal® susp 除癲達口服懸浮液	60 mg/ml, 100 ml/bot	4-16 yrs: 起始劑量: 8-10 mg/kg/day分2次 服用
* Valproate Sod. Depakine® solution 帝拔癲液	200 mg/ml, 40 ml/bot	20-30 mg/kg/day, BID - TID
<u>Misc. Anxiolytic, Sedatives, Hypnotics (其它鎮靜安眠劑)</u>		
Chloral hydrate 10% solution	2 mg/ml	Sedation: 25 mg/kg/day, Q8H Hypnotic: 50 mg/kg/dose
<u>GI Prokinetic Drugs (腸胃蠕動劑)</u>		
* Domperidone Wempty® susp 胃利空懸液劑	1 mg/ml, 60 ml/bot	0.25-0.5 ml/kg/dose TID-QID, before meal
<u>Cardiotonic Drugs (強心劑)</u>		
* Digoxin Cardiacin® elixir 地高新醣劑	50 mcg/ml, 60 ml/bot	10-20 mcg/kg/day
<u>Diuretics(利尿劑)</u>		
* Furosemide Fumide® oral solution 福滿內服液劑	10 mg/ml, 120 ml/bot	● 早產新生兒(<29 週): 1-2 mg/kg/dose Q24H ● 早產新生兒(>29 週): 1-2 mg/kg/dose Q12- 24H ● 新生兒: 1-3 mg/kg Q8H(需要時) ● 嬰兒及小孩: 起始劑量: 2 mg/kg/dose, 1-2 mg/kg/dose Q6-8H, MAX: 6mg/kg/dose
<u>Hormones & Synthetic substitutes- Adrenals(腎上腺素)</u>		
* Prednisolone Kidsolone® oral susp 必爾生口服液	1 mg/ml, 60 ml/bot	遵照醫囑
<u>Immunosuppressive Agents</u>		
* Cyclosporin Sandimmun® sol'n 新體睦口服溶液	100mg/ml, 50 ml/Bot	依各疾病適量投與，必要時依血中濃度調整劑量
<u>Iron Preparations (鐵製劑)</u>		
* Ferrum Hausmann® 富鐵好滴劑	50 mg iron /ml (1drop = 2.5 mg iron) 30 ml/bot	Infants: Begin with 6 drops daily and gradually increase to 20 drops daily Children: 20 drops 1-2 times daily

臺中榮總 兒科適用水劑藥品清單

藥品成份/商品名	成份含量	一般劑量
Replacement Preparations (取代用藥)		
Neutral Phosphate solution powder/pack	Phosphorus: 200 mg/10ml Sodium: 1.08 mEq/ml 500 ml/bot	視需要適量投與
Potassium Gluconate K-Glu®oral solution 15mL 鉀補口服液	鉀20 mEq /15mL/bot	<i>Neonatal:</i> 1 to 5 mEq/kg/day in divided doses; max single dose 2 mEq/kg. <i>Infants, Children, and Adolescents:</i> 2 to 5 mEq/kg/day in divided doses; max single dose 2 mEq/kg or 20 mEq.
Sodium Phosphate Dibasic Heptahydrate + Phosphoric Acid Phosphate oral solution 20mL 含磷口服液20毫升	磷30 mg/mL, 20mL/bot	2-3mL/kg/day(2-3mmol/kg/day)

- *有此星號標示之水劑為本院另有口服錠劑或膠囊劑型，因此對於較小的幼兒病患，請醫師們儘量開立水劑藥品。
- 滑鼠點選藥理分類標題即可超連結藥品手冊查詢系統。

藥學部 2023-05更新