## Baby D drop (Liquid form Vitamin D3 supplement, 15ml/Bot)

- Specially formulated for infants between the ages of 0-12 months
- Each 0.2ml of Baby D provides 5ug of Vitamin D3(cholecalciferol) and should be administered once per day (Do not exceed the stated dose)
- An oral syringe with a 0.2ml marking is provided in the pack to facilitate accurate dosing
- After use, the syringe can be cleaned with cooled boiled water
- Administer Baby D when your infant is in a relaxed state after feeding and winding
- Odorless and tasteless
- Use baby D within 3 months of opening

成分: 0.2ml = 5ug = 200IU (Vit.D: 1ug=40IU)

匯款資料:華南銀行 南永和分行

帳號 169100040363 優龍股份有限公司

單瓶價格 1200 元 打九折 1080 NTD

運費 100元 (\*\*兩瓶以上免運費)

請家屬匯款後,通知優龍股份有限公司出貨 02-25684868

寄至中榮加護病房或病嬰室,王主任或陳主任 收

\*\*中部一些藥局有鋪貨,要詢問看看;目前樓下維康沒有販售

營養標示(每 1ml 含有)

熱量:9.44 大卡

蛋白質: 0.001 公克

脂肪:1.05 公克

飽和脂肪 1.02 公克

反式脂肪 0 公克

碳水化合物 0 公克

鈉 0.004 毫克

維他命 D 25 微克



This leaflet gives you some useful information about using **Baby** D. Please read this leaflet carefully before using **Baby** D.

What is Baby 0?
Baby 0 is a Vitamin D, food supplement specially formulated for babies from birth onwards. Baby 0 comes in liquid form, in an amber glass bottle, along with a single oral dosing syringe.

Why use vitamin D<sub>3</sub>? Vitamin D<sub>3</sub> is a fat soluble vitamin. Vitamin D $_3$  is needed for normal growth and development of bone in children. The daily supplemented dose of Vitamin D $_3$  recommended for infants is 5  $\mu$ g (200 IU).

How should Baby D be given?
Baby D should be given orally.
Each 0.2 m Lof Baby D provides 5 µg (200 IU) of Vitamin D, (cholecalciferol) and should be administered once per day.

An oral syringe with a 0.2 mL marking is provided in the pack to facilitate accurate dosing. After use, the syringe can be cleaned with cooled boiled water.

Administer **Baby** D when your infant is in a relaxed state after feeding and winding. **Baby** D is odourless and tasteless making it easier for your baby to take.


Shake the bottle well before use. Remove the cap by pushing it down and turning it anticlockwise.

Remove the syringe from the wrapping, insert it into the opening at the neck of the bottle and dispose of the plastic wrapping immediately.

Turn the bottle upside down, draw back the white plunger of the syringe to the 0.2 mL mark. The tip of the plunger should be within the black outline as illustrated. Return the bottle to upright position and remove the syringe.

Place the open end of the syringe into your infant's mouth and press the plunger down slowly to deliver the contents. After use replace the cap back on the bottle, disassemble the syringe and wash parts in cooled boiled water.

As an alternative, **Baby** and can be administered on a sterilised spoon using the syringe provided to measure the dose.





Do not exceed the stated dose. If your baby misses a dose do not take a double dose. If you think you have overdosed contact your pharmacist or doctor.

Please note that high intakes of >25  $\mu$ g (1,000 IU) Vitamin D per day (which is 5 times the recommended daily allowance) can cause kidney and bone damage.

A food supplement is not a substitute for a healthy diet. Baby should be consumed as part of a varied and balanced diet and a healthy lifestyle.

Miglyol Oil Ph.Eur. Vitamin D<sub>3</sub> Ph.Eur. Antioxidant: Vitamin E

Baby D contains vitamin D, (cholecalciferol) formulated in miglyol oil.

Baby D contains trace amounts of Vitamin E, which act as an antioxidant for Vitamin D<sub>3</sub>.

All ingredients comply with Ph. Eur. (pharmaceutical standard). Miglyol oil is obtained from environmentally friendly sustainable resources.

Do not give **Baby** D if your baby is allergic to any of the ingredients.

Allergy Information

Baby 0 does not contain peanuts, peanut traces, gluten, lactose, sugars, crustaceans, eggs. fish, soya beans, celery, mustard or sesame seeds. Baby 0 does not contain oil from the following nuts almond, hazelnut, walnut, cashew, pecan nut, brazil nut, pistachio nut, macadamia nut or queensland nut.

Each pack contains 15 mL of **Baby D** in an amber glass bottle and an oral dosing syringe to administer the product. Should the syringe become damaged or lost please contact KoRa Healthcare for a replacement syringe

Storage
Keep Baby D and the pack contents out of the sight and reach of children

Store in a cool, dry place in original packaging and keep out of direct sunlight. Keep cap closed when not in use.

Do not use Baby 0 after the use by date, which is stated on the bottle and the carton.

Do not use Baby 0 if tamper evident seal has been broken before use.

Use Baby 0 within 3 months of opening. Each 15 mL bottle of Baby 0 contains 75 days supply.

oRa Healthcare vords Business Park, Swords, Co Dublin, Ireland eephone: 1800 42 62 82 ione: +353 (0) 1 8900406