

台中榮民總醫院  
內部過敏免疫風濕科

血管炎特別紀錄表(六)

Takayasu Arteritis

此處刷卡用

健保榮民	健保員工	健 保	健保員眷	民 眾	
姓名			病房	床位	年齡
					病歷號

**1990 criteria for the classification of Takayasu arteritis**

	Criteria	Definition
_____	1. Age at disease onset < 40 years	Development of symptoms or findings related to Takayasu arteritis at age $\leq$ 40 years
_____	2. Claudication of extremities	Development and worsening of fatigue and discomfort in muscles of 1 or more extremity while in use, especially the upper extremities
_____	3. Decreased brachial artery pulse	Decreased pulsation of 1 or both brachial arteries
_____	4. BP difference >10 mm Hg	Difference of >10 mm Hg in systolic blood pressure between arms
_____	5. Bruit over subclavian arteries or aorta	Bruit audible on auscultation over 1 or both subclavian arteries or abdominal aorta
_____	6. Arteriogram abnormality	Arteriographic narrowing or occlusion of the entire aorta, its primary branches, or large arteries in the proximal upper or lower extremities, not due to arteriosclerosis, fibromuscular dysplasia, or similar causes; changes usually focal or segmental

\* For purposes of classification, a patient shall be said to have Takayasu arteritis if at least 3 of these 6 criteria are present. The presence of any 3 or more criteria yields a sensitivity of 90.5% and a specificity of 97.8%. BP = blood pressure (systolic; difference between arms).

Date: \_\_\_\_\_ Signature: \_\_\_\_\_