

## 本院須作 penicillin 皮膚試驗之抗生素

| 抗生素類別                 | Penicillins   | Penicillins/ $\beta$ -Lactamase inhibitors   |
|-----------------------|---|--|
| 注射劑型<br>( Injection ) | <ul style="list-style-type: none"> <li>* <b>Ampicillin</b><br/>(Ampolin® inj 500 mg/vial)</li> <li>* <b>Benzathine Penicillin</b><br/>(Bicillin L-A® inj 2.4MU)</li> <li>* <b>Oxacillin</b><br/>(Ocillina® 500 mg/vial)</li> <li>* <b>Penicillin G</b><br/>(PCN G sod. 3MU/vial)</li> <li>* <b>Piperacillin</b><br/>(Piperacillin inj 2gm-CYH)</li> </ul> | <ul style="list-style-type: none"> <li>* <b>Amoxicillin/Clavulanate</b><br/>(Curam® 1.2 gm/vial,<br/>Soonmelt® inj 1.2g)</li> <li>* <b>Ampicillin/Sulbactam</b><br/>(Unasyn® 1.5 gm/vial)</li> <li>* <b>Piperacillin/Tazobactam</b><br/>(Tazocin® 2.25 gm/vial,<br/>Tapi® mycin Lyo 4.5 gm/vial<br/>Betamycin® inj 3.375Gm)</li> </ul> |

附註：根據護理部「盤尼西林皮膚試驗法」標準作業流程，一個月內有做過皮膚試驗者，可以不用再做。

### 以下口服劑型可以不用做皮膚試驗 但使用上還是要小心是否有任何過敏症狀

| 抗生素類別            | Penicillins   | Penicillins/ $\beta$ -Lactamase inhibitors  |
|------------------|---|---|
| 口服劑型<br>( Oral ) | <ul style="list-style-type: none"> <li>* <b>Amoxicillin</b><br/>(Supercillin® 250 mg/ cap )</li> <li>* <b>Ampicillin</b><br/>(Ampicillin 500mg/ cap)</li> <li>* <b>Dicloxacillin</b><br/>(Diclocin ® 250 mg/cap)</li> </ul> | <ul style="list-style-type: none"> <li>* <b>Amoxicillin/Clavulanate</b><br/>(Curam® Film-coated tab 1000mg, Curam® susp 312.5 mg/5 ml, Curam® dispersible tab 1000mg)</li> <li>* <b>Sultamicillin tosylate</b><br/>(Unasyn® 375mg/tab)</li> </ul> |