

臺中榮總 須作 penicillin 皮膚試驗之抗生素

抗生素類別	Penicillins	Penicillins/ β -Lactamase inhibitors
注射劑型 (Injection)	* Ampicillin (Ampolin[®] inj 500 mg/vial) * Benzathine Penicillin (Bicillin L-A[®] inj 2.4MU) * Oxacillin (Ocillina[®] 500 mg/vial) * Penicillin G (PCN G sod. 3MU/vial) * Piperacillin (Piperacillin inj 2gm-CYH)	* Amoxicillin/Clavulanate (Curam[®] 1.2 gm/vial, Soonmelt[®] inj 1.2g) * Ampicillin/Sulbactam (Unasyn[®] 1.5 gm/vial) * Piperacillin/Tazobactam (Tazocin[®] 2.25 gm/vial, Tapi[®] mycin Lyo 4.5 gm/vial, Betamycin[®] inj 3.375gm/vial)

以下口服劑型可以不用做皮膚試驗
但使用上還是要小心是否有任何過敏症狀

抗生素類別	Penicillins	Penicillins/ β -Lactamase inhibitors
口服劑型 (Oral)	* Amoxicillin (Supercillin[®] 250 mg/ cap, Amolin[®] oral susp 5% 60mL) * Ampicillin (Ampicillin 500mg/ cap) * Dicloxacillin (Diclocin[®] 250 mg/cap)	* Amoxicillin/Clavulanate (Curam[®] Film-coated tab 1000mg, Curam[®] susp 312.5 mg/5 ml, Curam[®] dispersible tab 1000mg) * Sultamicillin tosylate (Unasyn[®] 375mg/tab)