

台中榮民總醫院
內部過敏免疫風濕科

此處刷卡用

系統性紅斑狼瘡特別紀錄表

健保榮民	健保員工	健 保	健保員眷	民 眾	
姓名			病房	床位	年齡
					病歷號

The 1997 revised criteria for classification of SLE

- | Past | Now | |
|------|-----|---|
| () | () | 1. Malar rash :
Fixed erythema, flat or raised, over the malar eminences, tending to spare the nasolabial folds. |
| () | () | 2. Discoid rash :
Erythematous raised patches with adherent keratotic scaling and follicular plugging; atrophic scarring may occur in older lesions. |
| () | () | 3. Photosensitivity :
Rash as a result of unusual reaction to sunlight, by history , or physician observation. |
| () | () | 4. Oral ulcers :
Oral or nasopharyngeal ulceration, usually painless, observed by a physician. |
| () | () | 5. Arthritis :
Nonerosive arthritis involving 2 or more peripheral joints, characterized by tenderness, swelling or effusion. |
| () | () | 6. Serositis :
____(1)Pleuritis – convincing history of pleuritic pain or rub heard by a physician or evidence of pleural effusion; or
____(2)Pericarditis – documented by ECG or rubs or evidence of pericardial effusion |
| () | () | 7. Renal disorder :
____(1)Persistent proteinuria > 0.5g/day or > 3+, if quantitation not performed; or
____(2)Cellular casts –may be RBCs, hemoglobin, granular, tubular, or mixed. |
| () | () | 8. Neurologic disorder :
____(1)Seizure – in the absence of offending drugs, or known metabolic derangement, e.g., uremia, ketoacidosis, or electrolyte imbalance.
____(2)Psychosis – in the absence of offending drugs, or known metabolic derangement, Ketoacidosis, or electrolyte imbalance. |
| () | () | 9. Hematologic disorder :
____(1)Hemolytic anemia – with reticulocytosis ; or
____(2)Leukopenia < 4000/mm ³ total on 2 or more occasions; or
____(3)Lymphopenia < 1500/mm ³ on 2 or more occasions; or
____(4)Thrombocytopenia < 100,000/mm ³ in the absence of offending drugs |
| () | () | 10. Immunologic disorder :
____(1)Positive Antiphospholipid Antibody or
____(2)Anti - DNA : antibody to native DNA in abnormal titer; or
____(3)Anti - Sm : presence of antibody to Sm nuclear antigen; or
____(4)False – positive serologic test for syphilis known to be positive for at least 6 months and confirmed by Treponema pallidum immobilization or FTA-ABS. |
| () | () | 11. Antinuclear antibody (ANA) :
An abnormal titer of ANA by immunofluorescence or an equivalent assay at any point in time and in the absence of drugs known to be associated with “ drug-induced lupus ” syndrome. |