

台中榮民總醫院  
內部過敏免疫風濕科

血管炎特別紀錄表(二)

Wegener's Granulomatosis

此處刷卡用

健保榮民	健保員工	健 保	健保員眷	民 眾	
姓名			病房	床位	年齡
			病歷號		

**1990 Criteria for the Classification of Wegener's Granulomatosis**

	Criteria	Definition
_____	1. Nasal or oral inflammation	Development of painful or painless oral ulcers or purulent or bloody nasal discharge
_____	2. Abnormal chest radiograph	Chest radiograph showing the presence of nodules, fixed infiltrates, or cavities
_____	3. Urinary sediment	Microhematuria (>5 red blood cells per high power field) or red cell casts in urine sediment
_____	4. Granulomatous inflammation on biopsy	Histologic changes showing granulomatous inflammation within the wall of an artery or in the perivascular or extravascular area (artery or arteriole)

\* For purposes of classification, a patient shall be said to have Wegener's granulomatosis if at least 2 of these 4 criteria are present. The presence of any 2 or more criteria yields a sensitivity of 88.2% and a specificity of 92.0%

Date: \_\_\_\_\_ Signature: \_\_\_\_\_

*Leavitt RY, Fauci AS, Bloch DA, Michel BA, Hunder GG, Arend WP, et al. The American College of Rheumatology 1990 criteria for the classification of Wegener's granulomatosis. Arthritis Rheum 1990;33:1101---7.*

SAME6

©2007 American College of Rheumatology