

台中榮民總醫院
內部過敏免疫風濕科

此處刷卡用

混合性締結組織病特別紀錄表 (一)

健保榮民	健保員工	健 保	健保員眷	民 眾	
姓名			病房	床位	年齡
					病歷號

DIAGNOSTIC CRITERIA FOR MIXED CONNECTIVE TISSUE DISEASE

Alarcón-Segovia's Criteria ^[110]

- _____ 1. Serologic **criteria**
- _____ 2. Anti-RNP at hemagglutination
- _____ 3. B. Clinical **criteria**
- _____ 4. Swollen hands
- _____ 5. Synovitis
- _____ 6. Myositis (biologically proven)
- _____ 7. Raynaud's phenomenon
- _____ 8. Acrosclerosis
- _____ 9. Mixed connective tissue disorder (MCTD) present if:

Criterion A is accompanied by 3 or more clinical **criteria**, one of which must include synovitis or myositis

Kahn's Criteria ^[109]

- _____ 1. A. Serologic **criteria**
- _____ 2. High-titer anti-RNP corresponding titer of $\geq 1:1600$ to a speckled antinuclear antibodies (ANA) of $\geq 1:1200$ titer
- _____ 3. Clinical **criteria**
- _____ 4. Swollen fingers
- _____ 5. Synovitis
- _____ 6. Myositis
- _____ 7. Raynaud's phenomenon
- _____ 8. Mixed connective tissue disorder (MCTD) present if:

Criterion A is accompanied by Raynaud's phenomenon and two or more the three remaining clinical **criteria**