

台中榮民總醫院  
內部過敏免疫風濕科

此處刷卡用

系統性紅斑狼瘡特別紀錄表

健保榮民	健保員工	健 保	健保員眷	民 眾	
姓名			病房	床位	年齡
					病歷號

The 1982 revised criteria for classification of SLE

- |      |     |  |
|------|-----|--|
| Past | Now |  |
|------|-----|--|
- ( ) ( ) 1. **Malar rash** :  
Fixed erythema, flat or raised, over the malar eminences, tending to spare the nasolabial folds.
- ( ) ( ) 2. **Discoid rash** :  
Erythematous raised patches with adherent keratotic scaling and follicular plugging; atrophic scarring may occur in older lesions.
- ( ) ( ) 3. **Photosensitivity** :  
Rash as a result of unusual reaction to sunlight, by history, or physician observation.
- ( ) ( ) 4. **Oral ulcers** :  
Oral or nasopharyngeal ulceration, usually painless, observed by a physician.
- ( ) ( ) 5. **Arthritis** :  
Nonerosive arthritis involving 2 or more peripheral joints, characterized by tenderness, swelling or effusion.
- ( ) ( ) 6. **Serositis** :  
\_\_\_\_(1)Pleuritis – convincing history of pleuritic pain or rub heard by a physician or evidence of pleural effusion; or  
\_\_\_\_(2)Pericarditis – documented by ECG or rubs or evidence of pericardial effusion
- ( ) ( ) 7. **Renal disorder** :  
\_\_\_\_(1)Persistent proteinuria > 0.5g/day or > 3+, if quantitation not performed; or  
\_\_\_\_(2)Cellular casts – may be RBCs, hemoglobin, granular, tubular, or mixed.
- ( ) ( ) 8. **Neurologic disorder** :  
\_\_\_\_(1)Seizure – in the absence of offending drugs, or known metabolic derangement, e.g., uremia, ketoacidosis, or electrolyte imbalance.  
\_\_\_\_(2)Psychosis – in the absence of offending drugs, or known metabolic derangement, ketoacidosis, or electrolyte imbalance.
- ( ) ( ) 9. **Hematologic disorder** :  
\_\_\_\_(1)Hemolytic anemia – with reticulocytosis; or  
\_\_\_\_(2)Leukopenia < 4000/mm<sup>3</sup> total on 2 or more occasions; or  
\_\_\_\_(3)Lymphopenia < 1500/mm<sup>3</sup> on 2 or more occasions; or  
\_\_\_\_(4)Thrombocytopenia < 100,000/mm<sup>3</sup> in the absence of offending drugs
- ( ) ( ) 10. **Immunologic disorder** :  
\_\_\_\_(1)Positive LE cell preparation; or  
\_\_\_\_(2)Anti - DNA : antibody to native DNA in abnormal titer; or  
\_\_\_\_(3)Anti - Sm : presence of antibody to Sm nuclear antigen; or  
\_\_\_\_(4)False – positive serologic test for syphilis known to be positive for at least 6 months and confirmed by Treponema pallidum immobilization or FTA-Abs.
- ( ) ( ) 11. **Antinuclear antibody (ANA)** :  
An abnormal titer of ANA by immunofluorescence or an equivalent assay at any point in time and in the absence of drugs known to be associated with “ drug-induced lupus ” syndrome.