

台中榮民總醫院
內部過敏免疫風濕科

血管炎特別紀錄表(三)

Hypersensitivity vasculitis(1)

此處刷卡用

健保榮民	健保員工	健 保	健保員眷	民 眾	
姓名			病房	床位	年齡
					病歷號

1990 Criteria for the Classification of Hypersensitivity vasculitis

Criteria	Definition
_____ 1. Age at disease onset >16 years	Development of symptoms after age 16
_____ 2. Medication at disease onset	Medication was taken at the onset of symptoms that may have been a precipitating factor
_____ 3. Palpable purpura	Slightly elevated purpuric rash over one or more areas of the skin; does not blanch with pressure and is not related to thrombocytopenia
_____ 4. Maculopapular rash	Flat and raised lesions of various sizes over one or more areas of the skin
_____ 5. Biopsy including arteriole and venule	Histologic changes showing granulocytes in a perivascular or extravascular location

* For purposes of classification, a patient is said to have hypersensitivity vasculitis if at least 3 of these 5 criteria are present. The presence of any 3 or more criteria yields a sensitivity of 71.0% and a specificity of 83.9%.

+ This is criterion 20 from Table 1.

Date: _____ Signature: _____

Calabrese LH, Michel BA, Bloch DA, Arend WP, Edworthy SM, Fauci AS, et al. The American College of Rheumatology 1990 criteria for the classification of hypersensitivity vasculitis. Arthritis Rheum 1990;33:1108---13.

©2007 American College of Rheumatology